Author index

Adams, A.N., see Davies *et al.*, 400
Adams, Sally E., Jones, R.A.C. and Coutts, R.H.A. Effect of temperature on potato virus X infection in potato cultivars carrying different combinations of hypersensitivity genes, 517
Alexander, Helen Miller, Roelfs, A.P. and Cobbs, Gary. Effects of disease and plant competition on yield in monocultures and mixtures of two wheat cultivars, 457
Alonso, Angela, see Fraile *et al.*, 82
Andrews, Linda, see Locke, 241

Bailiss, K.W. and Ollennu, L.A.A. Effect of alfalfa mosaic virus isolates on forage yield of lucerne (*Medicago sativa*) in Britain, 162
Bailiss, K.W., see Omar *et al.*, 86; 535
Ballinger, D.J. and Kollmorgen, J.F. Control of take-all of wheat in the field with benzimidazole and triazole fungicides applied at seeding, 67
Ballinger, D.J. and Kollmorgen, J.F. Glasshouse and field evaluation of benomyl and triadimefon applied at seeding to control take-all in wheat, 61
Bar-Joseph, M., see Rosner *et al.*, 178
Barrett, J.A. Testing for heterothallism and mating-type in plant pathogenic fungi, 158
Ben-Yephet, Y., Bitton, S. and Greenberger, A. Control of lettuce drop disease, caused by *Sclerotinia sclerotiorum*, with metham-sodium soil treatment and foliar application of benomyl, 146
Bielenin, Anna, see Harris, 565
Billing, Eve, see Byrde *et al.*, 417
Bitton, S., see Ben-Yephet *et al.*, 146
Black, D.L.R. and Brown, Averil E. *Microdochium bolleyi* associated with root rot of flax, 592
Bonman, J.M., see Yeh, 319
Brasier, C.M., see Rogers *et al.*, 277
Brooke, R.C., see Melakeberhan *et al.*, 203
Brown, Averil, E., see Black, 592
Bruzzese, E. and Hasan, S. Infection of Australian and New Zealand *Rubus* subgenera *Dalibarda* and *Lampobatus* by the European blackberry rust fungus *Phragmidium violaceum*, 413
Buck, K.W., see Rogers *et al.*, 277
Butt, D.J. & Jeger, M.J. Components of spore production in apple powdery mildew (*Podosphaera leucotricha*), 491
Butt, D.J., see Jeger, 498; see Jeger *et al.*, 477
Byrde, R.J.W., Hutcheon, J.A., Williams, R.R., Billing, Eve and Gwynne, D.C. Fireblight at Long Ashton: a case history, 1971–1983, 417
Byrde, R.J.W., see Hutcheon *et al.*, 249

Café-Filho, Adalberto C., see Reifschneider *et al.*, 451
Callow, J.A., see Parry, 380; see Robinson, 169
Carpenter, J.M., see Katis *et al.*, 152
Carver, T.L.W. Histology of infection by *Erysiphe graminis* f. sp. *hordei* in spring barley lines with various levels of partial resistance, 232
Caten, C.E., see Newton *et al.*, 108; see Osbourn *et al.*, 135
Chapman, G.P., see Omar *et al.*, 86; 535
Cobbs, Gary, see Alexander *et al.*, 457
Cole, J.S., see Deall, 74
Coley-Smith, J.R. A comparison of flavour and odour compounds of onion, leek, garlic and *Allium fistulosum* in relation to germination of sclerotia of *Sclerotium cepivorum*, 370
Coley-Smith, J.R. Interactions between *Sclerotium cepivorum* and cultivars of onion, leek, garlic and *Allium fistulosum*, 362

Author index

Adams, A.N., see Davies *et al.*, 400
Adams, Sally E., Jones, R.A.C. and Coutts, R.H.A. Effect of temperature on potato virus X infection in potato cultivars carrying different combinations of hypersensitivity genes, 517
Alexander, Helen Miller, Roelfs, A.P. and Cobbs, Gary. Effects of disease and plant competition on yield in monocultures and mixtures of two wheat cultivars, 457
Alonso, Angela, see Fraile *et al.*, 82
Andrews, Linda, see Locke, 241

Bailiss, K.W. and Ollennu, L.A.A. Effect of alfalfa mosaic virus isolates on forage yield of lucerne (*Medicago sativa*) in Britain, 162
Bailiss, K.W., see Omar *et al.*, 86; 535
Ballinger, D.J. and Kollmorgen, J.F. Control of take-all of wheat in the field with benzimidazole and triazole fungicides applied at seeding, 67
Ballinger, D.J. and Kollmorgen, J.F. Glasshouse and field evaluation of benomyl and triadimefon applied at seeding to control take-all in wheat, 61
Bar-Joseph, M., see Rosner *et al.*, 178
Barrett, J.A. Testing for heterothallism and mating-type in plant pathogenic fungi, 158
Ben-Yephet, Y., Bitton, S. and Greenberger, A. Control of lettuce drop disease, caused by *Sclerotinia sclerotiorum*, with metham-sodium soil treatment and foliar application of benomyl, 146
Bielenin, Anna, see Harris, 565
Billing, Eve, see Byrde *et al.*, 417
Bitton, S., see Ben-Yephet *et al.*, 146
Black, D.L.R. and Brown, Averil E. *Microdochium bolleyi* associated with root rot of flax, 592
Bonman, J.M., see Yeh, 319
Brasier, C.M., see Rogers *et al.*, 277
Brooke, R.C., see Melakeberhan *et al.*, 203
Brown, Averil, E., see Black, 592
Bruzzese, E. and Hasan, S. Infection of Australian and New Zealand *Rubus* subgenera *Dalibarda* and *Lampobatus* by the European blackberry rust fungus *Phragmidium violaceum*, 413
Buck, K.W., see Rogers *et al.*, 277
Butt, D.J. & Jeger, M.J. Components of spore production in apple powdery mildew (*Podosphaera leucotricha*), 491
Butt, D.J., see Jeger, 498; see Jeger *et al.*, 477
Byrde, R.J.W., Hutcheon, J.A., Williams, R.R., Billing, Eve and Gwynne, D.C. Fireblight at Long Ashton: a case history, 1971–1983, 417
Byrde, R.J.W., see Hutcheon *et al.*, 249

Café-Filho, Adalberto C., see Reifschneider *et al.*, 451
Callow, J.A., see Parry, 380; see Robinson, 169
Carpenter, J.M., see Katis *et al.*, 152
Carver, T.L.W. Histology of infection by *Erysiphe graminis* f. sp. *hordei* in spring barley lines with various levels of partial resistance, 232
Caten, C.E., see Newton *et al.*, 108; see Osbourn *et al.*, 135
Chapman, G.P., see Omar *et al.*, 86; 535
Cobbs, Gary, see Alexander *et al.*, 457
Cole, J.S., see Deall, 74
Coley-Smith, J.R. A comparison of flavour and odour compounds of onion, leek, garlic and *Allium fistulosum* in relation to germination of sclerotia of *Sclerotium cepivorum*, 370
Coley-Smith, J.R. Interactions between *Sclerotium cepivorum* and cultivars of onion, leek, garlic and *Allium fistulosum*, 362

Coley-Smith, J.R., see Wang, 544; see Wang *et al.*, 427; see Wareing *et al.*, 390
Cook, R.J., see Royle *et al.*, 466
Cother, E.J. and Dowling, V. Bacteria associated with internal breakdown of onion bulbs and their possible role in disease expression, 329
Coutts, R.H.A., see Adams *et al.*, 517; see Henry *et al.*, 585
Cowland, Janet A., see Hooper, 128
Coyle, J., see Hutcheon *et al.*, 249
Cox, T.W., see Mitchell *et al.*, 403
Crute, I.R. Investigations of gene-for-gene relationships: the need for genetic analyses of both host and parasite, 15

Damgaci, E., see Hewett, 377
Darby, P., Lewis, B.G. and Matthews, P. Diversity of virulence within *Ascochyta pisi* and resistance in the genus *Pisum*, 214
D'Arcy, C.J. and Hewings, A.D. Enzyme-linked immunosorbent assays for study of serological relationships and detection of three luteoviruses, 288
Davies, D.L., Stickels, J.E. and Adams, A.N. A single occurrence of apple proliferation disease in England, 400
Day, P.R. Plant–parasite interaction: a genetical perspective, 262
Deall, M.W. and Cole, J.S. A comparative study of the pathogenicity and epidemiology of strains of *Pseudomonas syringae* pv. *tabaci* that cause wildfire and angular leaf spot diseases of tobacco in Zimbabwe, 74
Deverall, B.J., see Loughman, 443
Dixon, G.R. and Robinson, D.L. The susceptibility of *Brassica oleracea* cultivars to *Plasmodiophora brassicae* (clubroot), 101
Dowling, V., see Cother, 329
Dugard, P.I., see Kennedy *et al.*, 344
Duncan, J.M., see Kennedy *et al.*, 344

Elphinstone, J.G. and Pérombelon, M.C.M. Contamination of potatoes by *Erwinia carotovora* during grading, 25

Fahy, P.C., see Moffett *et al.*, 34
Fraile, Aurora, Alonso, Angela and Sagasta, E.M. Some characteristics of *Botrytis cinerea* isolates tolerant to procymidone, 82
Francki, R.I.B., Ryan, C.C., Hatta, T., Rohozinski, J. and Grivell, C.J. Serological detection of Fiji disease virus antigens in the planthopper *Perkinsiella saccharicida* and its inefficient ability to transmit the virus, 324
Fyfe, A.M., see Tantius *et al.*, 578

Garrett, Constance M.E. Influence of rootstock on the susceptibility of sweet cherry scions to bacterial canker, caused by *Pseudomonas syringae* pvs *morsprunorum* and *syringae*, 114
Gaunt, R.E., see Lim, 44; 54
Gibson, R.W., see Katis *et al.*, 152
Ginzburg, I., see Rosner *et al.*, 178
Gregory, P.H. Obituary, 134
Gregory, R.S., see Hollins *et al.*, 93
Greenberger, A., see Ben-Yephet *et al.*, 146
Grivell, C.J., see Francki *et al.*, 324
Guile, C.T., see Port, 575
Gutteridge, R.A., see Plumb *et al.*, 314
Gwynne, D.C., see Byrde *et al.*, 417

Hamadi, A., see Martelli, 595

Hammond, Kim E. and Lewis, B.G. The timing and sequence of events leading to stem canker disease in populations of *Brassica napus* var. *oleifera* in the field, 551
Harris, D.C. and Bielenin, Anna. Evaluation of selective media and bait methods for estimating *Phytophthora cactorum* in apple orchard soils, 565
Hasan, S., see Bruzzese, 413
Hatta, T., see Francki *et al.*, 324
Hattingh, M.J., see Mansvelt, 337
Hayward, A.C. *see* Moffett *et al.*, 34
Henry, Christone M., Jones, R.A.C. and Coutts, R.H.A. Occurrence of a soil-borne virus of sugar beet in England, 585
Hewett, P.D. and Damgaci, E. A new procedure to detect a low incidence of *Ustilago nuda* in seed barley, 377
Hewings, A.D., see D'Arcy, 288
Hide, G.A. *Phoma* and *Fusarium* rots on imported 'new' potatoes, 126
Hignett, R.C., Zamse, S.E., Munro, S.M. and Smith, A.R.W. Inhibition by bacterial lipopolysaccharide of wilt induced by phytopathogenic bacteria in cut cherry shoots, and of cherry canker disease, 270
Hiruki, C. A new yellows disease of *Baekea virgata* associated with mycoplasma-like organisms, 396
Hoare, F.A., Hunter, T. and Jordan, V.W.L. Influence of spray programmes on development of fungicide resistance in the eyespot pathogen of wheat, *Pseudocercosporella herpotrichoides*, 506
Holgate, M.E., see Hutcheon *et al.*, 249
Holliday, J.M., see Stone *et al.*, 18
Hollins, T.W., Scott, P.R. and Gregory, R.S. The relative resistance of wheat, rye and triticale to take-all caused by *Gaeumannomyces graminis*, 93
Hooper, D.J. and Cowland, Janet A. Fungal hosts for the chrysanthemum nematode, *Aphelenchoides ritzemabosi*, 128
Humpherson-Jones, F.M. The occurrence of virulent pathotypes of *Leptosphaeria maculans* in brassica seed crops in England, 224
Hunter, T., see Hoare *et al.*, 506
Hutcheon, J.A., Coyle, J., Holgate, M.E. and Byrde, R.J.W. Effects of fungicides on long-term cropping and fruit quality of apple, 249
Hutcheon, J.A., see Byrde *et al.*, 417

Irwin, J.A.G., see Stirling, 527

Jeger, M.J. Asymptotic behaviour and threshold criteria in model plant disease epidemics, 355
Jeger, M.J. and Butt, D.J. Epidemics of apple powdery mildew (*Podosphaera leucotricha*) in a mixed cultivar orchard, 498
Jeger, M.J., Butt, D.J. and Swait, A.A.J. Components of resistance of apple to powdery mildew (*Podosphaera leucotricha*), 477
Jeger, M.J., see Butt, 491
Jellis, G.J., see Mitchell *et al.*, 403
Johnson, R., see Newton, *et al.*, 108
Jones, R.A.C., see Adams *et al.*, 517; see Henry *et al.*, 585
Jordan, V.W.L., see Hoare *et al.*, 506

Katan, T., see Shabi *et al.*, 310
Katis, N., Carpenter, J.M. and Gibson, R.W. Interference between potyviruses during aphid transmission, 152
Kennedy, D.M., Duncan, J.M., Dugard, P.I. and Topham, P.H. Virulence and aggressiveness of single-zoospore isolates of *Phytophthora fragariae*, 344
Kenny, M.A., see Wainwright *et al.*, 582
Ko, W.H., Tomita, Judy and Short, R.L. Two natural hosts of *Kretzschmaria clavus* in Hawaiian forests, 254
Koenraadt, H., see Shabi *et al.*, 310

Kollmorgen, J.F., see Ballinger, 61; 67

Lennon, E.A., see Plumb *et al.*, 314
Lester, E. Objectives and perspectives in plant disease control, 2
Lewis, B.G., see Darby *et al.*, 214; see Hammond, 551
Lim, L.G. and Gaunt, R.E. The effect of powdery mildew (*Erysiphe graminis* f. sp. *hordei*) and leaf rust (*Puccinia hordei*) on spring barley in New Zealand. II. Apical development and yield potential, 54
Lim, L.G. and Gaunt, R.E. The effect of powdery mildew (*Erysiphe graminis* f. sp. *hordei*) and leaf rust (*Pucchinia hordei*) on spring barley in New Zealand. I. Epidemic development, green leaf area and yield, 44
Locke, T. and Andrews, Linda. Effects of fungicides on powdery mildew, tree growth and cropping of apple, 241
Loughman, R. and Deverall, B.J. Infection of resistant and susceptible cultivars of wheat by *Pyrenophora tritici-repentis*, 443
Lumb, V.M., Pérombelon, M.C.M. and Zutra, D. Studies of a wilt disease of the potato plant in Israel caused by *Erwinia chrysanthemi*, 196

Mansfield, J.W., see Omar *et al.*, 535
Mansvelt, E.L. and Hattingh, M.J. Pear blossom blast in South Africa caused by *Pseudomonas syringae* pv. *syringae*, 337
Martelli, G.P. and Hamadi, A. Occurrence of eggplant mottled dwarf virus in Algeria, 595
Mathias, P.L., see Stone *et al.*, 18
Matthews, P., see Darby *et al.*, 214
Mayoral, M.L., see Rosner *et al.*, 178
McNicol, J.W., see Phillips, 185
Melakeberhan, H., Brooke, R.C. and Webster, J.M. Relationship between physiological response of French bean of different age to *Meloidogyne incognita* and subsequent yield loss, 203
Mitchell, A.G., see Webber *et al.*, 512
Mitchell, S.J., Jellis, G.J. and Cox, T.W. *Sclerotinia sclerotiorum* on linseed, 403
Moffett, Melda, L., Hayward, A.C. and Fahy, P.C. Five new hosts of *Pseudomonas andropogonis* occurring in eastern Australia: host range and characterization of isolates, 34
Moore, F.J.H. Obituary, 259
Munro, S.M., see Hignett *et al.*, 270

Newton, A.C., Johnson, R. and Caten, C.E. Attempted somatic hybridization of *Puccinia striiformis* f. sp. *tritici* and *P. striiformis* f. sp. *hordei*, 108

Ollennu, L.A.A., see Bailiss, 162
Omar, S.A.M., Bailiss, K.W. and Chapman, G.P. Virus-induced changes in the responses of faba bean to infection by *Botrytis*, 86
Omar, S.A.M., Bailiss, K.W., Chapman, G.P. and Mansfield, J.W. Effects of virus infection of faba bean on subsequent infection by *Uromyces viciae-fabae*, 535
Osbourn, A.E., Scott, P.R. and Caten, C.E. The effects of host passaging on the adaptation of *Septoria nodorum* to wheat or barley, 135

Parrott, D.M., see Stone *et al.*, 18
Parry, R.W.H. and Callow. J.A. The dynamics of homologous and heterologous interactions between rice and strains of *Xanthomonas campestris*, 380
Pérombelon, M.C.M., see Elphinstone, 25; see Lumb *et al.*, 196
Phillips, M.S. and McNicol, J.W. The use of biplots as an aid to interpreting interactions between potato clones and populations of potato cyst nematodes, 185
Plumb, R.T., Lennon, E.A. and Gutteridge, R.A. The effects of infection by barley yellow mosaic virus on the yield and components of yield of barley, 314

Port, C.M. and Guile, C.T. Outbreaks of flea beetles, *Altica* spp., on heather and other flowering plants, 575

Raccah, B., see Rosner *et al.*, 178
Rego, Arildo M., see Reifschneider *et al.*, 451
Reifschneider, Francisco J.B., Café-Filho, Adalberto C. and Rego, Arildo M. Factors affecting expression of resistance in pepper (*Capsicum annuum*) to blight caused by *Phytophthora capsici* in screening trials, 451
Robinson, J.N. and Callow, J.A. Multiplication and spread of pathovars of *Xanthomonas campestris* in host and non-host plants, 169
Robinson, D.L., see Dixon, 101
Roelfs, A.P., see Alexander *et al.*, 457
Rogers, H.J., Buck, K.W. and Brasier, C.M. Transmission of double-stranded RNA and a disease factor in *Ophiostoma ulmi*, 277
Rohozinski, J., see Francki *et al.*, 324
Rosner, A., Raccah, B., Mayoral, M.L., Bar-Joseph, M. and Ginzburg, I. Synthesis of DNA complementary to the polyadenylated genomic RNA of potato virus Y and its molecular cloning, 178
Royle, D.J., Shaw, M.W. and Cook, R.J. Patterns of development of *Septoria nodorum* and *S. tritici* in some winter wheat crops in Western Europe, 1981–83, 466
Rudgard, S.A. Witches' broom disease on cocoa in Rondonia, Brazil: basidiocarp production on detached brooms in the field, 434
Ryan, C.C., see Francki *et al.*, 324

Sagasta, E.M., see Fraile *et al.*, 82
Scott, P.R., see Hollins *et al.*, 93; see Osbourn *et al.*, 135
Shabi, E., Koenraadt, H. and Katan, T. Further studies of the inheritance of benomyl resistance in *Venturia pirina* isolated from pear orchards in Israel, 310
Shattock, R.C., see Tantius *et al.*, 578; see Thomas, 120
Shaw, D.S., see Tantius *et al.*, 578
Shaw, M.W. Effects of temperature and leaf wetness on *Pyrenophora teres* growing on barley cv. Sonja, 294
Short, R.L., see Ko *et al.*, 254
Shaw, M.W., see Royle *et al.*, 466
Smith, A.R.W., see Hignett *et al.*, 270
Smith, Frances, see Webber *et al.*, 512
Stickels, J.E., see Davies *et al.*, 400
Stirling, A.M. and Irwin, J.A.G. Etiology of a newly described root rot of guar (*Cyamopsis tetragonoloba*) in Australia caused by *Phytophthora cryptogea*, 527
Stone, A.R., Holliday, J.M., Mathias, P.L. and Parrott, D.M. A selective survey of potato cyst-nematode pathotypes in Great Britain, 18
Swait, A.A.J., see Jeger *et al.*, 477

Tantius, P.H., Fyfe, A.M., Shaw, D.S. and Shattock, R.C. Occurrence of the A2 mating type and self-fertile isolates of *Phytophthora infestans* in England and Wales, 578
Thomas, M.R. and Shattock, R.C. Effects of fungicides on *Drechslera* spp. and leaf surface filamentous saprophytic fungi on *Lolium perenne*, 120
Tomita, Judy, see Ko *et al.*, 254
Topham, P.H., see Kennedy *et al.*, 344

Wainwright, A., Kenny, M.A. and Yarham, D.J. Increased susceptibility to *Septoria nodorum* in winter wheat infected by *Tilletia caries*, 582
Wang, Zu-Nan and Coley-Smith, J.R. Studies on some characteristics of dicarboximide-resistant isolates of *Botrytis cinerea* from protected lettuce, 544

Wang, Zu-Nan, Coley-Smith, J.R. and Wareing, P.W. Dicarboximide resistance in *Botrytis cinerea* in protected lettuce, 427
Wang, Z.-N., see Wareing *et al.*, 390
Wareing, P.W., Wang, Z.-N. and Coley-Smith, J.R. Fungal pathogens in rotted basal leaves of lettuce in Humberside and Lancashire with particular reference to *Rhizoctonia solani*, 390
Wareing, P.W., see Wang *et al.*, 427
Webber, Joan, Mitchell, A.G. and Smith, Frances. Linkage of the genes determining mating type and fungicide tolerance in *Ophiostoma ulmi*, 512
Webster, J.M., see Melakeberhan *et al.*, 203
Williams, R.R., see Byrde *et al.*, 417

Yarham, D.J., see Wainwright *et al.*, 582
Yeh, W.H. and Bonman, J.M. Assessment of partial resistance to *Pyricularia oryzae* in six rice cultivars, 319

Zamse, S.E., see Hignett *et al.*, 270
Zutra, D., see Lumb *et al.*, 196